

Kink Dynamics in a Lattice Model with Long-Range Interactions

T. Ioannidou^{1*}, J. Pouget¹ and E. Aifantis^{2†}

¹*Laboratoire de Modélisation en Mécanique (associé au CNRS),
Université Pierre et Marie Curie, Tour 66, 4 place Jussieu,
75252 Paris Cédex 05, France*

²*Laboratory of Mechanics and Materials, Polytechnic School,
Aristotle University of Thessaloniki, 54006, Thessaloniki, Greece*

Abstract

This paper proposes a one-dimensional lattice model with long-range interactions which, in the continuum, keeps its nonlocal behaviour. In fact, the long-time evolution of the localized waves is governed by an asymptotic equation of the Benjamin-Ono type and allows the explicit construction of moving kinks on the lattice. The long-range particle interaction coefficients on the lattice are determined by the Benjamin-Ono equation.

1 Introduction

This paper is concerned with the effects of long range interactions in the behaviour of solitons (kinks) in a lattice system. In particular, we show that the introduced lattice model asymptotically leads to a nonlocal continuum equation. By nonlocal we mean the model where the localized solution is of order of the scale parameter (*ie* parameter which has the dimension of length), and thus it is physically acceptable to consider wavelengths comparable with the scale parameter. Its equation of motion contains integral, integro-differential or finite difference operators in the spatial variables, while the wave propagation velocity depends on the wavelength. In the present model there are two characteristic lengths: (a) the lattice spacing and (b) the radius of particle interaction range.

The nonlocal models can be divided into two classes: discrete and continuous. For specific cases, it is simpler and easier to use the quasicontinuum description of the discrete

*Permanent Address: Institute of Mathematics, University of Kent, Canterbury CT2 7NF, UK

†Center for Mechanics of Materials and Instabilities, Michigan Technological University, Houghton, MI 49931, USA

medium. Its essence is an interpolation of functions of discrete argument by a special class of analytic functions in such a way that a correspondence condition between the quasicontinuum and the discrete medium is fulfilled. The advantages of such an approach is the description of discrete and continuum media within a unified formalism and their correct generalization. In that case the quasicontinuum model is also applicable to macrosystems. Analytic solutions valid for wide nonlinear excitations have been obtained using the continuum approximation in [1, 2] and the quasicontinuum approximation in [3, 4, 5]. In particular in [1, 6], it has been shown that very narrow soliton-like excitations can propagate without any energy loss due to discreteness effects.

The lattice model we study, possesses an inherent nonlocal behaviour, *ie* the motion of a particle at node n depends on the motion of its neighbours. Such models for different kinds of particle interactions have been studied in a series of papers [7, 8, 9]. So far the classical procedure for studying them, at the continuum limit, is to assume slowly varying fields and expand them using Taylor's series. Then, approximate partial differential equations for the continuous fields were obtained.

In this paper, we show how to keep the nonlocal behaviour of the lattice model at the continuum level, or in other words, how to describe the nonlocality at the continuum level. This transition from the continuum to the discrete system with long-range interactions and vice-versa, is straightforward due to the quasicontinuum approximation. The idea of preserving the nonlocality at the continuum limit occurs in condensed-matter physics to describe phase transition in crystals, nonlinear waves in crystals and biological molecules. For example, modified versions of the discrete nonlinear Schrödinger equation which keep the nonlocality at the continuum and describe problems in biomolecules have been studied in the past [10, 11].

It is worthwhile mentioning some interesting studies of anharmonic atomic chain including long-range interactions due to large Coulomb coupling between particles [12, 13, 14, 15]. The long-range potential is of the inverse power-type or Kac-Baker form. In the case of inverse power type of the fourth-order, the continuum limit yields an integro-differential equation involving a Hilbert transform [15]. Using a perturbative technique the authors derive a mixed modified Korteweg-de-Vries with Benjamin-Ono equations [15], which is close to the result presented in the following.

The paper is organised as follows: In Section 2 we introduce the lattice model; while in Section 3, we deal with its continuum approximation and show that it leads to the Benjamin-Ono equation for long-time evolution. In Section 4, we derive the type of long-range particle interactions on the lattice using the fact that the lattice model asymptotically leads to the Benjamin-Ono equation. Finally, numerical simulations of the discrete model based on the Benjamin-Ono soliton are given and discussed in Section 5.

where \sum'_n means summation over all $n \neq 0$. This equation is deduced from the lattice energy invariance; *ie* the invariance by translation of the chain as a whole.

Assume now, that the particles interact nonlocally with measure $\Psi(n)$. Then it can be shown that [16]

$$\Phi(n) = -\Psi(n), \quad n \neq 0; \quad \Phi(0) = \sum'_n \Psi(n). \quad (2.4)$$

Note that for real mechanical systems all $\Psi(n) > 0$ and hence $\Phi(0) > 0$, $\Phi(n) < 0$ ($n \neq 0$). However, a system may be constructed, for which some particles would have negative interaction coefficients; which is the case of our model described below.

3 The Continuum Approximation

The set of nonlinear differential equations for a discrete system is generally complex and, quite often, its continuum approximation is considered (which is the case here). By assuming that the displacements are slowly varying over the lattice spacing, the continuum approximation may be derived as follows:

3.1 The Nonlocal Model

The Lagrangian (2.1) in the form which is invariant with respect to n and x representations, can be written as

$$\mathcal{L} = \frac{1}{2} \langle \rho \dot{u}, \dot{u} \rangle + \frac{1}{2} \langle u, \Phi u \rangle - \frac{1}{3} \langle \Delta u, k_3 \Delta u^2 \rangle, \quad (3.5)$$

where ρ is the mass density, while $u = u_n(t)$, $\Delta u = b^{-1}(u_{n+1} - u_n)$ and $u = u(x, t)$, $\Delta u = u_x$ in the discrete and continuum space.

The various terms in (3.5) for both the discrete and the continuum representations are given as follows

$$\begin{aligned} \frac{1}{2} \langle u, \Phi u \rangle &= \frac{1}{4} \sum_{nn'} \Phi(n - n') (u_n - u_{n'})^2 \\ &= \frac{1}{2} \int_{R^2} \Phi(x - x') u(x) u(x') dx dx', \end{aligned} \quad (3.6)$$

$$\begin{aligned} \frac{1}{3} \langle \Delta u, k_3 \Delta u^2 \rangle &= \frac{1}{3} \sum_n k_3 (u_{n+1} - u_n)^3 \\ &= \frac{1}{3} \int_R k_3 u_x^3 dx. \end{aligned} \quad (3.7)$$

The above forms contain elastic (long-range) interactions for both the discrete model and its quasicontinuum counterparts.

The equation of motion of the medium in the (x, t) representation is

$$\rho \ddot{u}(x, t) = - \int_R \Phi(x - x') u(x', t) dx' + k_3 (u_x^2)_x. \quad (3.8)$$

The corresponding condition (2.3) in the x representation becomes $\int_R \Phi(x - x') dx' = 0$ which implies that $\Phi(x - x')$ can be represented as

$$\Phi(x - x') = \psi(x) \delta(x - x') - \Psi(x - x'), \quad (3.9)$$

with $\Psi(x - x') = \Psi(x' - x)$ and $\psi(x) \doteq \int \Psi(x - x') dx'$.

Its quasicontinuum counterparts are of the form

$$\Phi(n - n') = \psi(n) \delta(n - n') - \Psi(n - n'), \quad (3.10)$$

with $\Psi(n - n') = \Psi(n' - n)$ and $\psi(n) \doteq \sum_{n'} \Psi(n - n')$. Thus $\Psi(n - n')$ and $\Psi(x - x')$ are the interaction coefficients and the interaction coefficient density of the model connecting the points (n, n') and (x, x') , respectively.

In the x representation the force constant can be expressed as [16]

$$\Phi(x - x') = \frac{\partial^2}{\partial x \partial x'} c(x - x'), \quad (3.11)$$

where $c(x - x')$ is the so-called kernel operator of elastic moduli.

Then, equation (3.8) after integration by parts becomes

$$\rho \ddot{u}(x, t) = \frac{\partial}{\partial x} \int_R c(x - x') \frac{\partial u(x')}{\partial x'} dx' + k_3 (u_x^2)_x, \quad (3.12)$$

which is quite similar to the equation obtained from the nonlocal elasticity of the one-dimensional linear case (*ie* with the k_3 term vanishing) [17, 18].

In this case, by using the kinematical definition for the strain $\epsilon(x', t) = \partial u(x', t) / \partial x'$ and the nonlocal constitutive equation for the stress $\sigma(x, t) = \int_{-l}^l c(x - x') \epsilon(x', t) dx'$ where l is the interaction range length, equation (3.12) can be recast into

$$\rho \ddot{u}(x, t) = \sigma_x, \quad (3.13)$$

which is the equation of motion of a one-dimensional elastic medium. [Note that, the integral over a finite radius of interaction defines the finiteness of action at a distance determined by this radius (that is, $l = \mathcal{M}b$). Specifically, $\Phi(y)$ is not zero for $y \in [-l, l]$ and it equals to zero otherwise; while, if the chain is bounded within an interval $2L$ then $l \ll L$.]

In this connection, equation (3.13) is the starting point in continuum mechanics and expresses the balance of linear momentum. It is endowed with appropriate constitutive equations for the stress leading to partial differential equations for the determination of the displacement field. For example, a simple linear elastic model of the form $\sigma = E\epsilon$ with E denoting the Young modulus (identified as c_0^2 below) leads to the classical wave equation

$u_{tt} = c^2 u_{xx}$ ($c^2 \equiv E/\rho$). If gradient-dependent nonlinear elasticity model is adopted [19] of the form $\sigma = f(\epsilon) - \tilde{c}\epsilon_{xx}$ the appropriate equation of motion reads $f(u_x)u_{xx} - \tilde{c}u_{xxx} = \rho u_{tt}$ which contain the $k_3(u_x^2)_x$ term listed above, even though a more interesting case results when $f'(u_x)$ is non-monotonic. Other types of such constitutive models and the corresponding equations of motion are discussed in [20]. A case of particular interest in terms of its discrete model counterpart arises when viscosity is considered.

For a simple Kelvin-Voigt model of the form $\sigma = E\epsilon + \mu\dot{\epsilon}$ where μ is the viscosity and $\dot{\epsilon}$ the strain rate, the resultant equation of motion with $\lambda \equiv \mu/\rho$ is $u_{tt} = c^2 u_{xx} + \lambda \dot{u}_{xx}$, *ie* the damped wave equation. It is an open question to examine a Lagrangian-like lattice version of this model.

On the other hand, by letting

$$c = c_0^2 \delta(x - x') + c_1(x - x'), \quad (3.14)$$

where c corresponds to the local response with c_0^2 denoting the local elastic modulus and c_1 describing the nonlocal behaviour of the model, equation (3.12) simplifies to

$$\rho \ddot{u}(x, t) = c_0^2 u_{xx} + \frac{\partial}{\partial x} \int_R c_1(x - x') \frac{\partial u(x')}{\partial x'} dx' + k_3(u_x^2)_x, \quad (3.15)$$

which is further examined below.

3.2 The Long-Time Evolution of the Localized Wave

In this section we investigate the asymptotic behaviour of equation (3.15) for large times. By assuming that the contribution of the nonlinear term is significant throughout we rescale the nonlinear and the dispersive term by introducing a small parameter $\varepsilon < 1$, as follows

$$2k_3 = \gamma \varepsilon, \quad c_1 = \varepsilon g. \quad (3.16)$$

Here, γ is the rescaled nonlinearity coefficient and g is the rescaled elastic kernel. Then, for large times, the asymptotic expansion for the displacement field can be expressed as

$$u(x, t) = u_0(\xi, \tau) + \varepsilon u_1(x, t) + O(\varepsilon^2), \quad (3.17)$$

where $\xi = x - vt$ is a shifted coordinate and $\tau = \varepsilon t$ is the slow time variable.

Using equations (3.16) and (3.17) and keeping terms of order ε only, equation (3.15) becomes

$$2\rho v u_{0\xi\tau} + \frac{\partial}{\partial \xi} \int_R g(\xi - \xi') u_{0\xi'} d\xi' + \gamma u_{0\xi} u_{0\xi\xi} = -c_0^2 u_{1xx} + \rho u_{1tt}, \quad (3.18)$$

where $c_0^2 = \rho v^2$.

Then, the secularity condition [21] implies that both the left and right-hand sides of (3.18) are zero. Thus, the right-hand side leads to the standard linear wave equation for u_1 ; while the left-hand side gives the long-time behaviour of u_0 , *ie*

$$2\rho v u_{0\xi\tau} + \frac{\partial}{\partial \xi} \int_R g(\xi - \xi') u_{0\xi'} d\xi' + \gamma u_{0\xi} u_{0\xi\xi} = 0. \quad (3.19)$$

Finally, by letting

$$u_{0\xi} = \mathcal{F}, \quad g(z) = \frac{\partial}{\partial z} G(z), \quad (3.20)$$

where $z = \xi - \xi'$ and $y = \xi - z$, equation (3.19) becomes

$$2\rho v \mathcal{F}_\tau + \gamma \mathcal{F} \mathcal{F}_\xi + \int_R G(\xi - y) \frac{\partial^2}{\partial^2 y} \mathcal{F}(y, \tau) dy = 0, \quad (3.21)$$

which transforms to the Benjamin-Ono equation for

$$\tau = \frac{1}{2\rho v} T, \quad \mathcal{F} = \frac{1}{\gamma} U, \quad G(\xi - y) = -\frac{\alpha}{\pi} \frac{1}{\xi - y}. \quad (3.22)$$

Note that, under the swap of the role played by x and t in (3.21), we arrive at an integro-differential equation usually met in nonlinear wave propagation in active media with dissipative process [22]. The latter is due to viscoelasticity described by an integral law involving exponential kernel in the integral part of the equation of motion. This kernel is then a function of time introduced by a relaxation process.

3.3 The Benjamin-Ono Equation

The Benjamin-Ono evolution equation

$$U_T + U U_\xi + \alpha H(U_{\xi\xi}) = 0, \quad (3.23)$$

with α being a positive parameter, was originally derived in [23, 24, 25] for interval-wave propagation in a two-layer system: one shallow and the other infinitely deep. In this context, $U(\xi, T)$ represents the amplitude of the interfacial wave produced by an initial disturbance, say $U(\xi, 0)$. The operator H designates the one-dimensional Hilbert transform, *ie*

$$\alpha H(U_{\xi\xi}) = \frac{\alpha}{\pi} \int_R \frac{1}{y - \xi} U_{yy} dy, \quad (3.24)$$

which implies that (3.23) is an integro-differential weakly nonlinear evolution equation.

Equation (3.23) has a simple solitary wave solution in the form of a Lorentzian (algebraic) shape [21, 26], *ie*

$$U(\xi, T) = \frac{4\nu}{(\frac{\nu}{\alpha})^2 (\xi - \nu T)^2 + 1}, \quad (3.25)$$

where 4ν is the amplitude of the wave and $\frac{\alpha}{\nu}$ measures the wavelength. The velocity of the soliton is amplitude-dependent and equals to ν .

Remark: Regarding the long-wavelength limit of the lattice model, the classical continuum approximation (*ie* Taylor's expansion for the fields) of (3.15) leads to the local nonlinear partial derivative equation of the Boussinesq type [1, 2]

$$u_{tt} - c^2 u_{xx} + a u_x u_{xx} + b u_{xxx} = 0, \quad (3.26)$$

where the parameters c , a and b depend on the atomic interactions. Alternatively, equation (3.26) may be directly deduced as a special case from the continuum approximation of (2.2).

The long-time evolution of (3.26) is given by the Korteweg-de-Vries equation,

$$V_t + VV_x + V_{xxx} = 0, \quad (3.27)$$

when multi-scale techniques (or perturbation methods) as described in Section 3.2 are used. In addition, the Benjamin-Ono equation (3.23) can be transformed into the Korteweg-de-Vries equation by replacing $\alpha H(U_{\xi\xi})$ by $U_{\xi\xi\xi}$; however, this transition is not an obvious one.

4 Evaluation of the Interaction Coefficients

In this section we derive the lattice interaction coefficients $\Psi(n)$ of the model (2.1) using the fact that its asymptotic expansion, for large times, leads to the Benjamin-Ono equation; and therefore, the kernel of the operator of elastic moduli $c(x)$ is known.

Since there is a correspondence between functions of discrete argument and a certain class of analytic functions, the interaction coefficients can be evaluated either in the space of functions of discrete argument or in the space of analytic functions. However, since all these spaces are isomorphic to each other the interaction coefficients are obtained using the x representation (due to simplicity).

Equation (3.11) by setting $x - x' \rightarrow x$ becomes

$$\begin{aligned} \Phi(x) &= -\frac{\partial^2}{\partial x^2} c(x) \\ &= -c_0^2 \delta''(x) - \varepsilon g''(x), \end{aligned} \quad (4.28)$$

where (3.14) and (3.16) were used. But from the long-time evolution analysis of the localized wave the explicit form of the function g is known. Thus, using (3.20) and (3.22) we get

$$\begin{aligned} \Phi(x) &= -c_0^2 \delta''(x) - \varepsilon G'''(x) \\ &= -c_0^2 \delta''(x) - \frac{6\varepsilon\alpha}{\pi} \frac{1}{x^4}. \end{aligned} \quad (4.29)$$

Due to Fourier transform, the force constants in the space of discrete argument are related with the ones in the space of analytic functions as [16]:

$$\begin{aligned} \Phi(n) = \Phi(nb) &= \int_R \Phi(x) \delta_b(x - nb) dx \\ &= -c_0^2 \mathcal{I}_1(n) - \frac{6\alpha\varepsilon}{\pi} \mathcal{I}_2(n), \end{aligned} \quad (4.30)$$

where $\delta_B(x) \doteq (2\pi)^{-1} \int_B e^{ikx} dx = \sin(\pi x)(\pi x)^{-1}$ and

$$\begin{aligned}\mathcal{I}_1(n) &= \int_R \delta''(x) \frac{\sin[\frac{\pi}{b}(x-nb)]}{\pi(x-nb)} dx, \\ \mathcal{I}_2(n) &= \int_R \frac{1}{x^4} \frac{\sin[\frac{\pi}{b}(x-nb)]}{\pi(x-nb)} dx.\end{aligned}\tag{4.31}$$

Thus, it is a matter of algebra to evaluate the two integrals and obtain

$$\begin{aligned}\mathcal{I}_1(0) &= -\frac{\pi^2}{3b^3}, \\ \mathcal{I}_2(0) &= \frac{\pi^4}{24b^4},\end{aligned}\tag{4.32}$$

while for $n \neq 0$

$$\begin{aligned}\mathcal{I}_1(n) &= -\frac{2}{b^3} \frac{(-1)^n}{n^2}, \\ \mathcal{I}_2(n) &= \frac{\pi^2}{b^4} \frac{(-1)^n}{n^2} \left[\frac{1}{2} + \frac{(-1)^n - 1}{n^2 \pi^2} \right].\end{aligned}\tag{4.33}$$

Therefore, the lattice force constants are given by

$$\begin{aligned}\Phi(0) &= \frac{c_0^2 \pi^2}{3b^3} - \frac{\alpha \varepsilon \pi^3}{4b^4}, \\ \Phi(n) &= \frac{2c_0^2}{b^3} \frac{(-1)^n}{n^2} - \frac{6\alpha \varepsilon \pi}{b^4} \frac{(-1)^n}{n^2} \left[\frac{1}{2} + \frac{(-1)^n - 1}{n^2 \pi^2} \right], \quad n \neq 0.\end{aligned}\tag{4.34}$$

Finally, the lattice interaction coefficients $\Psi(n)$ in (2.2) (due to (3.10)) are given by

$$\Psi(n) = \Phi(0)\delta(n) - \Phi(n).\tag{4.35}$$

As one can observe from Figure 2, by keeping the nonlocality of the discrete model to the continuum one, and specifically, the choice of an integrable nonlocal equation as its asymptotic limit for large times, leads to the construction of a lattice model with negative interaction coefficients between its particles. In fact, Figure 2 illustrates the change of sign of $\Psi(n)$ and its variation with n for an atomic chain where each particle interacts with its ten neighbour particles. [The values of the other parameters, *ie* (a, b, ε, c_0) are listed in the figure caption.]

5 Numerical Simulations

The behaviour of genuinely discrete systems is quite different than the one of their continuum counterparts and these differences have been studied in numerous papers both from




Figure 2: The function $\Psi(n)$ given by (4.34)-(4.35) for $\alpha = 1$, $\nu = 1/4$, $\varepsilon = 0.1$, $n_0 = 125$ and phase velocity (a) $c_0 = 0.6$ solid line (b) $c_0 = 0.65$ dashed line (c) $c_0 = 0.7$ dotted line.

the mathematical and the physical point of view. Many of the methods used and their physical implementation in models with nearest neighbour interactions can be found in [27, 28, 29]. In addition, in [30, 31] the effects of discreteness in solitary waves in nearest neighbours lattices have been mathematically investigated.

In our case, the lattice model keeps its nonlocal behaviour at the continuum when its asymptotical limit for large times is the Benjamin-Ono equation. However, it is not obvious if the solution of the Benjamin-Ono equation will be the solution of the lattice model, as well. Thus, in this section we investigate the dynamical behaviour of the lattice model (2.2) numerically, using as an initial condition the soliton solution given in Section 3.3.

We propose a numerical scheme by directly considering the lattice equations (2.2), employing a Runge-Kutta method of fourth order and imposing pseudo-periodic boundary conditions, *ie*

$$u(N + i) = u(i) + u_0, \quad i \in [-\mathcal{M}, \mathcal{M}], \quad (5.36)$$

where N is the number of particles in the lattice, \mathcal{M} is the range of particle interactions considered and u_0 is the amplitude of the initial kink.

The initial conditions for the displacements and the velocities of each lattice particle are given by the analytic expression

$$u(x, t) = \frac{2\varepsilon\alpha}{k_3} \arctan\left[\frac{\nu}{\alpha} \left(x - x_0 - c_0 \left(1 + \frac{\varepsilon\nu}{2c_0^2}\right) t\right)\right], \quad (5.37)$$

and its time derivative, respectively. Here x_0 denotes the soliton position and we have set the mass density equal to one ($\rho = 1$). This solution represents a kink with thickness $\Delta = \alpha\pi/\nu$, *ie* the number of particles participating in the kink solution is proportional to




Figure 3: (a) Initial soliton solution (3.25) and (b) Initial kink solution (5.37) for $\alpha = 1$, $\nu = 1/4$ and $n_0 = 125$.

the kernel factor α and decreases as the wave parameter ν increases. [The solution (5.37) has been obtained from the solitary wave solution of the Benjamin-Ono equation (3.25) after using the change of variables and fields given in Section 3.2]. Figure 3 illustrates (a) the soliton solution $U(\zeta)$ with $\zeta = \xi - \nu T$ given by (3.25) and (b) the kink solution $u(x)$ given by (5.37) for the choice $b = 1$.

In all our simulations we have chosen the following fixed values for: $k_3 = 0.1$, $\varepsilon = 0.1$, $b = 1$, $\alpha = 1$, $M = 1$. However, different values for the above parameters give qualitatively the same results. The initial position of the kink is chosen equal to $n_0 = 125$, while the value of the kink velocity c_0 is chosen so that the first interaction coefficient is positive (*ie* $\Psi(1) > 0$) and the total number of particles in the lattice is equal to 250.

Figure 4 summarises the main results of our numerical simulations when models with different number of non-nearest neighbour interactions are considered and different values for the velocity and wavelength of the initial kink have been chosen. In particular, Figures 4(a), 4(e) and 4(f) present a solution corresponding to a lattice model with first, tenth and twentieth neighbour particle interactions when the initial kink wavelength and phase velocity are: $\nu = 1/8$, $c_0 = 0.7$; $\nu = 1/4$, $c_0 = 0.6$; and $\nu = 1/8$, $c_0 = 0.6$, respectively. In each model, the kink contains from twelve up to twenty four particles, which means that the continuum approximation is valid. [Recall that the kink thickness depends on the parameter ν , due to (5.37)]. On the other hand, Figures 4(b), 4(c) and 4(d) present the kink dynamics for lattice models with second, fourth and seventh neighbour particle interactions. However, we choose the kink width in all three cases to be small, *ie* the kink contains six particles with $\nu = 1/2$, $c_0 = 0.54$ in the first case, and three particles with $\nu = 1$; $c_0 = 0.6$, $c_0 = 0.7$ in the last two.

From Figure 4, one can observe that the initial soliton solution of the Benjamin-Ono equation is also a solution of the lattice model with long-range interactions given by (2.2). In fact, the initial lattice kink relaxes and propagates in the lattice space with small oscillations. It can be observed from Figures 4(a), 4(e) and 4(f) that as the kink thickness increases (*ie* ν decreases) the lattice radiation is eliminated. Numerical simulations of the

discrete nonlocal model for kink widths equal to three or six lattice spacings have been presented in Figures 4(b)-4(d): at this level the kink changes slightly and loses energy by radiation of small amplitudes waves. In all cases where the kink contains at least 20 particles, it is remarkable stable and emits (almost) no radiation.

Let us conclude by saying that, in the long-wavelength approximation (ie, when the average kink wavelength is larger compared to the lattice spacing) the initial (continuum) kink, which has been derived using the quasi-continuum approximation of the discrete model, relaxes to a stable discrete kink without any oscillations and our method is accurate.

6 Concluding Remarks

The present work is concerned with the dynamics of lattice models with long-range and nonlinear interactions. The main objective is the description of the dynamics of the lattice model, at the continuum level, by keeping the nonlocal nature of the discrete system. Our model is the simplest one consisting of a one-dimensional chain of point masses connected by springs. We have developed a theory for such a discrete system and introduced the notion of quasicontinuum which allows us to treat the discrete and continuum models in the framework of the same formalism. Attention has been focused to the analysis of an approximate model and its transition to nonlocal elasticity in the limit of long waves. In this formalism, we have set up a one-to-one correspondence between functions of discrete arguments and a class of analytic functions. The procedure is also applicable for integral operations on these functions. The method brings out a resemblance to the problem of interpolation of a discrete function by a smooth function satisfying some smoothness conditions. The technique allows us (Section 3) to use the same representation of the Lagrangian for all cases: discrete and analytic.

The equation of motion of the model in the continuum representation is similar to that of a nonlocal model of elasticity including a local nonlinear term. Nevertheless, such an equation of motion is not easy to deal with and a perturbation method has been employed. More precisely, we have considered the long-time evolution of nonlinear waves traveling in the lattice. The asymptotic method has been used to transform the equation of motion into an equation of the Benjamin-Ono type which governs the long-time evolution of the initial signal. For a particular choice of the nonlocal elastic kernel or interaction force function, the dispersive operator of the equation is a Hilbert transform corresponding to the Benjamin-Ono equation. In this particular situation, a localized wave solution of soliton type has been obtained. In order to test the analytical conjectures thus developed, some numerical simulations have been performed directly on the lattice model for different ranges of nonlocal action. It has been observed from numerics that the initial soliton (kink) solution propagates almost without perturbation and it is very stable. Moreover, wide kinks propagate in the lattice without any loss due to discreteness radiations.

Such a study can be extended to other kinds of nonlinear terms (double-well or sine-potential, higher-order interactions) or to arbitrary particle interactions. Extension to two-dimensional lattices can also be envisaged. The problems of kink collision or the influence




Figure 4: Kink plane for (a) one-range interaction, $\nu = 1/8$, $c_0 = 0.7$ (b) two-range interaction, $\nu = 1/2$, $c_0 = 0.54$ (c) four-range interaction, $\nu = 1$, $c_0 = 0.6$ (d) seven-range interaction, $\nu = 1$, $c_0 = 0.7$ (e) ten-range interaction, $\nu = 1/4$, $c_0 = 0.6$ (f) twenty-range interaction, $\nu = 1/8$, $c_0 = 0.6$.

of perturbations (forces, damping, lattice defects) on the kink motion is of interest. Some of these problems will be addressed in the future.

Acknowledgements

This work has been performed in the framework of the TMR European Contract **FMRX-CT-960062**: *Spatio-temporal instabilities in deformation and fracture mechanics, material science and nonlinear physics aspects*. TI acknowledges the Nuffield Foundation for a newly appointed lecturer award.

References

- [1] Peyrard M., Pnevmatikos St. and Flytzanis N.: Discreteness effect on non-topological kink soliton dynamics in nonlinear lattices, *Physica D* **19**, 268-281 (1986).
- [2] Flytzanis N., Pnevmatikos St. and Remoissenet M.: Soliton resonances in atomic nonlinear systems, *Physica D* **26**, 311-320 (1987).
- [3] Collins M.A.: Some properties of large amplitude motion in an anharmonic chain with nearest neighbour interactions, *J. Chem. Phys.* **77**, 2607-2622 (1986).
- [4] Roseneau Ph.: Dynamics of dense lattices, *Phys. Rev. B* **36**, 5868-5876 (1987).
- [5] Roseneau Ph.: Quasi-continuum spatial motion of a mass spring chain, *Physica D* **27**, 224-234 (1987).
- [6] Hochstrasser D., Mertens F.G. and Buttner H.: An iterative method for calculation of narrow solitary excitations on atomic chains, *Physica D* **35**, 259-266 (1989).
- [7] Askar A.: *Lattice Dynamical Foundations of Continuum Theories: Elasticity, Piezoelectricity, Viscoelasticity, Plasticity*, Series in Theoretical and Applied Mechanics, Vol. **2** (World Scientific Publishing Co.,Singapore, 1985).
- [8] Remoissenet M. and Flytzanis N.: Solitons in anharmonic chains with long-range interactions, *J. Phys. C: Solid State Phys.* **18**, 1573-1584 (1985).
- [9] Maradudin A.A., Montrol E.M., Weiss G.H. and Ipotova I.P.: *Theory of Lattice Dynamics in the Harmonic Approximation*, Solid State Physics, Suppl. 3 (academic Press, New York, 1971).
- [10] Johansson M., Gaididei Y., Christiansen P. and Rasmussen K.: Switching between states in a discrete nonlinear model with long-range dispersion, *Phys. Rev. E* **57**, 4739-4742 (1998).

- [11] Mingaleev S., Christiansen P., Gaididei Y., Johansson M. and Rasmussen K.: Models for energy and charge transport and storage in biomolecules, *J. Biol. Phys.* **25**, 41-63 1999
- [12] Kosevich A.M. and Kovalev A.S.: Self-localization of vibrations in a one-dimensional anharmonic chain, *Sov. Phys. JETP* **40**, 891-896 1975
- [13] Gaididei Y., Flytzanis N., Nueper A. and Mertens F.G.: Effects on nonlocal interactions on soliton dynamics in anharmonic chains: Scale competition, *Physica D* **107**, 83-111 1997
- [14] Bonart D., Rössler T., and Page J.B.: Intrinsic localized modes in complex lattice dynamical systems, *Physica D* **113**, 123-128 1998
- [15] Carstea A.S., Grecu D. and Visinescu A.: Exact solutions of a mixed KDV + mKDV and Benjamin-Ono equation, *Phys. Lett. A* **246**, 82-86 1998
- [16] Kunin I.A.: *Elastic Media with Microstructure I: One Dimensional Models*, Springer Series in Solid-State Sciences Vol. **26** (Springer - Verlag, Berlin, 1982).
- [17] Kröner E.: Elasticity theory of materials with long-range cohesive forces, *Int. J. Solids Structures* **3**, 731-742 (1967).
- [18] Eringen A.C.: Linear theory of nonlocal elasticity and dispersion of plane waves, *Int. J. Engng. Sci.* **10**, 425-435 (1972).
- [19] Triantafyllidis N. and Aifantis E.C.: A gradient approach to localization of deformation. 1-Hyperelastic materials, *J. Elasticity* **16**, 225-237 (1986).
- [20] Aifantis E.C.: Non-linearity, periodicity and patterning in plasticity and fracture, *Int. J. Non-linear Mech.* **31**, 797-809 (1996).
- [21] Nayfeh A.: *Perturbation Methods*, (John Wiley & Sons, New York, 1973).
- [22] Fridman V.E.: Steady waves in active birelaxating media, *Research Reports in Physics*, Ed. J. Engelbrecht (Springel-Verlag, Berlin, 1989) 24-51.
- [23] Benjamin T.B.: Internal waves of permanent form in fluid of great depths, *J. Fluid. Mech.* **29**, 559-592 (1967).
- [24] Davis R.E. and Acrivos A.: Solitary internal waves in deep water, *J. Fluid. Mech.* **29**, 593-607 (1967).
- [25] Ono H.: Algebraic solitary waves in stratified fluids, *J. Phys. Soc. Japan* **39**, 1082-1091 (1975).
- [26] Abdelouhab L., Bona J.L., Felland M. and Saut J.C.: Nonlocal models for nonlinear dispersive waves, *Physica D* **40**, 360-392 (1989).

- [27] Balmforth N., Craster R. and Kevrekidis P.: Being stable and discrete, *Physica D* **135**, 212-232 (2000).
- [28] Kevrekidis P., Jones C.K.R.T: Bifurcation of internal solitary wave modes from the essential spectrum, *Phys. Rev. E* **61**, 3114-3121 (2000).
- [29] Kevrekidis P., Jones C.K.R.T., Kapitula T.: Exponentially small splitting of heteroclinic orbits: from the rapidly forced pendulum to discrete solitons, *Phys. Lett. A* **269**, 120-129 (2000).
- [30] Kevrekidis P. and Weinstein M.: Dynamics of lattice kinks, *Physica D* **142**, 113-152 (2000).
- [31] Johansson M. and Aubry S.: Growth and decay of discrete nonlinear Schrodinger breathers interacting with internal modes or standing-wave phonons, *Phys. Rev. E* **61**, 5864-5879 (2000).